


2ND ORDER PARTIAL DERIVATIVES



The only thing more fun than differentiating once
is differentiating twice!

The only thing more fun than differentiating once
is differentiating twice!

$$z = x^2 - y^2$$

The only thing more fun than differentiating once is differentiating twice!

$$z = x^2 - y^2$$

$$z_x = \frac{\partial z}{\partial x} = 2x$$

$$z_y = \frac{\partial z}{\partial y} = -2y$$

The only thing more fun than differentiating once is differentiating twice!

$$z = x^2 - y^2$$

$$z_x = \frac{\partial z}{\partial x} = 2x$$

$$z_y = \frac{\partial z}{\partial y} = -2y$$

$$z_{xx} = (z_x)_x = \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = 2$$

The only thing more fun than differentiating once is differentiating twice!

$$z = x^2 - y^2$$

$$z_x = \frac{\partial z}{\partial x} = 2x$$

$$z_y = \frac{\partial z}{\partial y} = -2y$$

$$z_{xx} = (z_x)_x = \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = 2$$

$$z_{yy} = (z_y)_y = \frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial y} \right) = -2$$

The only thing more fun than differentiating once is differentiating twice!

$$z = x^2 - y^2$$

$$z_x = \frac{\partial z}{\partial x} = 2x$$

$$z_y = \frac{\partial z}{\partial y} = -2y$$

$$z_{xx} = (z_x)_x = \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = 2$$

$$z_{yy} = (z_y)_y = \frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial y} \right) = -2$$

$$z_{xy} = (z_x)_y = \frac{\partial^2 z}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial x} \right) = 0$$

The only thing more fun than differentiating once is differentiating twice!

$$z = x^2 - y^2$$

$$z_{xx} = (z_x)_x = \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = 2$$

$$z_{yy} = (z_y)_y = \frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial y} \right) = -2$$

$$z_{xy} = (z_x)_y = \frac{\partial^2 z}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial x} \right) = 0$$

$$z_{yx} = (z_y)_x = \frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial y} \right) = 0$$

$$z_x = \frac{\partial z}{\partial x} = 2x$$

$$z_y = \frac{\partial z}{\partial y} = -2y$$

It's no accident in this example that
the mixed partials are equal, $z_{xy} = z_{yx}$.

It's no accident in this example that
the mixed partials are equal, $z_{xy} = z_{yx}$.

This is what happens most of the time.

It's no accident in this example that the mixed partials are equal, $z_{xy} = z_{yx}$. This is what happens most of the time.

Theorem: If z_{xy} and z_{yx} are continuous at (a,b) , and interior point of the domain, then $z_{xy}(a,b) = z_{yx}(a,b)$.

What do the 2nd order partials tell us?

$$z = x^2 - y^2$$

$$z_x = 2x$$

$$z_y = -2y$$

$$z_{xx} = 2$$

$$z_{yy} = -2$$

In this case, they tell us that for a fixed value of y , the curve of intersection will be concave up.


$$z = x^2 - y^2$$

$$z_x = 2x$$

$$z_y = -2y$$


$$z_{xx} = 2$$

$$z_{yy} = -2$$



And for a fixed value of x ,
the curve of intersection will be concave down.

$$z = x^2 - y^2$$
$$z_x = 2x \qquad z_{xx} = 2$$
$$z_y = -2y \qquad z_{yy} = -2$$



The geometric interpretations of the mixed partials are harder to visualize. However, the good news is that we're not going to have to worry about them for what we want to accomplish.

$$z = x^2 - y^2$$

$$z_x = 2x$$

$$z_{xy} = 0$$

$$z_y = -2y$$

$$z_{yx} = 0$$

