

DIVERGENCE THEOREM IN THREE DIMENSIONS

In each problem below you are given a volume V bounded by a surface S along with a vector field F . If N is an outward pointing unit normal vector, then the outward flux across the surface created by the vector field F is, by the Divergence (Gauss') Theorem, equal to $\iint_S \vec{F} \cdot N dS = \iiint_V \operatorname{div} \vec{F} dV = \iiint_V \nabla \cdot F dV$. Evaluate this integral for each problem below

1. V is the solid ball with surface S defined by $x^2 + y^2 + z^2 = 1$

$$F = x\hat{i} + y\hat{j} + z\hat{k}$$

2. V is the cube defined by $0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1$, and S is the surface of the cube

$$F = x^2\hat{i} + y\hat{j} + z\hat{k}$$

V is the cylinder defined by $-1 \leq x \leq 1, -\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}, 0 \leq z \leq 4$,

3. and S is the surface of the cylinder

$$F = y\hat{i} - x\hat{j}$$

V is the solid bounded by the xy -plane and the hemisphere $z = \sqrt{4-x^2-y^2}$,

4. and S is the surface of V .

$$F = x^3\hat{i} + y^3\hat{j} + z^3\hat{k}$$

V is the solid bounded by the portion in the first octant of the

5. cylinder defined by $x^2 + y^2 = 1$ as z varies from 0 to 1,
and S is the surface of V .

$$F = x^2\hat{i}$$