

STOKES' THEOREM IN THREE DIMENSIONS

In each problem below you are given a surface S , defined by $z = f(x, y)$, over a region R , defined by the given limits on x and y . Let C_R be the boundary of the region R , oriented counterclockwise, and let C be the corresponding bounding curve on the surface S , also oriented counterclockwise (Except for problem 4. On problem 4, let your bounding curves be oriented clockwise.). Then if F is a vector field and N is the upward pointing unit normal vector for the surface S , use the higher dimensional version of Stokes' Theorem, $\int_C \vec{F} \cdot d\vec{r} = \iint_S (\text{curl } F \cdot N) dS$, to measure the circulation around the curve C that is caused by the vector field F .

$$S: z = -x^2 - y^2 + 4$$

1. $R: 0 \leq x \leq 1, 0 \leq y \leq 2$

$$F = z\hat{i} + x\hat{j} + y\hat{k}$$

$$S: z = y$$

2. $R: 0 \leq x \leq 1, 0 \leq y \leq 1 - x$

$$F = -3y^2\hat{i} + 4z\hat{j} + 6x\hat{k}$$

$$S: z = x^2 - y^2$$

3. $R: -1 \leq x \leq 1, -\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}$

$$F = x^2\hat{i} + y^2\hat{j} + z^2\hat{k}$$

$$S: z = 1 - x^2 - y^2$$

4. $R: -1 \leq x \leq 1, -\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}$

$$F = z\hat{i} + x\hat{j} + y\hat{k}$$

(NOTE: On this problem, let C_R and C be oriented clockwise. This means that your unit normal N will be pointing downward instead of upward.)

$$S: z = \frac{1}{2}\sqrt{1-x^2-y^2}$$

5. $R: -1 \leq x \leq 1, -\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}$

$$F = x^2\hat{i} + y^2\hat{j} + z^2 \tan xy\hat{k}$$

(HINT: Use a simpler surface with the same bounding curve.)