


# The Second Partial Test



**Definition:** Let  $(a,b)$  be a point contained in an open region  $R$  on which a function  $z = f(x, y)$  is defined. Then  $(a,b)$  is a **critical point** if any of the following conditions are true:

1.  $z_x(a,b) = 0 = z_y(a,b)$
2.  $z_x(a,b)$  does not exist
3.  $z_y(a,b)$  does not exist

**Definition:** Let  $(a,b)$  be a point contained in an open region  $R$  on which a function  $z = f(x, y)$  is defined. Then  $(a,b)$  is a **critical point** if any of the following conditions are true:

1.  $z_x(a,b) = 0 = z_y(a,b)$
2.  $z_x(a,b)$  does not exist
3.  $z_y(a,b)$  does not exist

**Theorem:** If  $z = f(x, y)$  has a local maximum or a local minimum at a point  $(a,b)$  contained within an open region  $R$  on which  $z = f(x, y)$  is defined, then  $(a,b)$  is a **critical point**.

**Second Partial Test:** Suppose  $z = f(x, y)$  has continuous second partial derivatives on an open region containing a point  $(a, b)$  such that

$z_x(a, b) = 0 = z_y(a, b)$ , and let

$$D = D(a, b) = \begin{vmatrix} z_{xx}(a, b) & z_{xy}(a, b) \\ z_{yx}(a, b) & z_{yy}(a, b) \end{vmatrix} = z_{xx}(a, b)z_{yy}(a, b) - z_{xy}(a, b)z_{yx}(a, b).$$

**Second Partial Test:** Suppose  $z = f(x, y)$  has continuous second partial derivatives on an open region containing a point  $(a, b)$  such that


$z_x(a, b) = 0 = z_y(a, b)$ , and let

$$D = D(a, b) = \begin{vmatrix} z_{xx}(a, b) & z_{xy}(a, b) \\ z_{yx}(a, b) & z_{yy}(a, b) \end{vmatrix} = z_{xx}(a, b)z_{yy}(a, b) - z_{xy}(a, b)z_{yx}(a, b).$$

Then:

1. If  $D > 0$  and  $z_{xx}(a, b) > 0$ ,  $f(a, b)$  is a **local minimum**.
2. If  $D > 0$  and  $z_{xx}(a, b) < 0$ ,  $f(a, b)$  is a **local maximum**.
3. If  $D < 0$ ,  $(a, b, f(a, b))$  is a **saddle point**.
4. If  $D = 0$ , the test is **inconclusive**.

Example:  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$



**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$



**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

critical point



**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

critical point



$$z_{xx} = 4 \quad z_{xy} = 2$$

$$z_{yx} = 2 \quad z_{yy} = 2$$

**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

critical point



$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

$$z_{xx} = 4 \quad z_{xy} = 2$$

$$z_{yx} = 2 \quad z_{yy} = 2$$

$$D(-1,1) = \begin{vmatrix} 4 & 2 \\ 2 & 2 \end{vmatrix} = 8 - 4 = 4 > 0$$

$$z_{xx}(-1,1) = 4 > 0$$

**Example:**  $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

critical point




$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

$$\begin{array}{ll} z_{xx} = 4 & z_{xy} = 2 \\ z_{yx} = 2 & z_{yy} = 2 \end{array} \quad D(-1,1) = \begin{vmatrix} 4 & 2 \\ 2 & 2 \end{vmatrix} = 8 - 4 = 4 > 0$$
$$z_{xx}(-1,1) = 4 > 0$$

Therefore,  $f(-1,1) = -4$  is a **local minimum**,  
and  $(-1,1,-4)$  is a **minimum point**.

Try it now with  $z = f(x, y) = x^3 - 3x + y^3 - 3y$ !



Also try  $z = f(x, y) = x^4 - y^4$ ,  $z = f(x, y) = x^4 + y^4$ , and  $z = -x^4 - y^4$ .

